

附件一

Table 2C. *Staphylococcus* spp. (Continued)

Antimicrobial Agent	<i>Staphylococcus</i> spp. Indications	Disk Content	Interpretive Categories and Zone Diameter Breakpoints, nearest whole mm				Interpretive Categories and MIC Breakpoints, µg/mL			
			S	SDD	I	R	S	SDD	I	R
Linezolid	All staphylococci	30 µg	≥ 26	–	23–25	≤ 22	≤ 4	–	–	≥ 8

<i>Staphylococcus</i> spp.							
Antibiotics	Version	Disk diffusion (mm)			MIC (µg/mL)		
		S	I	R	S	I	R
Linezolid	CLSI M100 S34 新標準	≥ 26	23-25	≤ 22	≤ 4	-	≥ 8
	CLSI M100 S33 舊標準	≥ 21	-	≤ 20	≤ 4		≥ 8

附件二

Table 2B-4. *Stenotrophomonas maltophilia* (Continued)

Antimicrobial Agent	Disk Content	Interpretive Categories and Zone Diameter Breakpoints, nearest whole mm			Interpretive Categories and MIC Breakpoints, µg/mL		
		S	I	R	S	I	R
TETRACYCLINES							
Minocycline	30 µg	≥ 26	21–25	≤ 20	≤ 1	2	≥ 4

<i>Stenotrophomonas maltophilia</i>							
Antibiotics	Version	Disk diffusion (mm)			MIC (µg/mL)		
		S	I	R	S	I	R
Minocycline	CLSI M100 S34 新標準	≥ 26	21-25	≤ 20	≤ 1	2	≥ 4
	CLSI M100 S33 舊標準	≥ 19	15-18	≤ 14	≤ 4	8	≥ 16