



Prevention of Radiation Dermatitis – Barrier Film Spray 放射性皮膚炎之預防 噴灑保膚膜注意事項(英文)

No Sting Barrier Film is a high-molecular polymer solution that forms a layer of transparent, breathable and waterproof protective film with lasting effect like a second layer of skin. It does not contain alcohol, is painless and can effectively isolate the irritation from urine, stool, secretion, and scotch tape.

No Sting Barrier Film can reduce the radiation dermatitis caused by radiation therapy. Follow medical instruction for the usage and time. Use it approximately once every 2 ~ 3 days.

Procedures

- Before using the No Sting Barrier Film on the area, clean the area and allow it to completely dry before using this product.
- First pull the overlapping and wrinkle area of skin, then spray it on the body. Keep the nozzle in 10 ~ 15 cm away from the skin, and at 40 degree angle, evenly spray the film on the affected area. Let the product dry for 30 seconds so the spray can form the protective film, before releasing the skin wrinkles. To reapply or spray in some area, allow the first layer of No Sting Barrier Film to dry (approximately 30 0seconds) before applying the next layer. Place the cap back to the nozzle after use to prevent spray by accident.

Precautions

- Skin susceptible to friction such as outer breast, intergluteal cleft of hips, armpit, and groin, use once in every 2 days.
- Wet are such as perineum, use once a day.
- For smooth and thin skin such as the face, neck and area with poor blood supply, use once in every 3 days.

- The product will naturally detach from the skin due to metabolism in 72 hours, which will not reside. Do not rub the skin to remove.
- The normal cleaned and sprayed area will not disappear due to bathing. Do not use the product with cream, lotion and ointment.
- External use only.

若有任何疑問，請不吝與我們聯絡
電話：(04) 22052121 分機 7450
HE-RA006-1-E